

the dry specimen. Flowers yellowish-red. Sepals subequal, broadly ovate. Petals slightly smaller than sepals, cuneately obovate. Lip 3-lobed with bottle-shaped spur at the base, apex of the spur globose, didynamous; side lobes rounded and midlobe oblong. Stigmatic surface 1 x 1 mm. Capsule not seen.

Fl.: Late June-late July

Fr.: Unknown.

Geographical distribution: India, Assam, Sikkim, Bangladesh, Bhutan and Thailand.

Specimen examined: Chittagong Hill

Tracts (Ragamati); Kaptai, 10.ii.1981 (collected at vegetative stage, subsequently flowered next year, cultured in the Botanical garden, Chittagong University, Chittagong, Bangladesh), Moyeen 404.

April 22, 1998

MOKTER AHMED

M.K. PASHA

Department of Botany,
University of Chittagong,
Chittagong 4331,
Bangladesh.

REFERENCES

- AIRY-SHAW, H.K. (Ed.) 1973: A Dictionary of the Flowering Plants and Ferns by J.C. Willis (Ed.) 8. Cambridge University Press, England. pp 1001.
- BLUME, C.L. (1825): Bijdragen tot. de Fl. Van Nedert. Indie-Batavia. p. 434.
- BOSE, T.K. & S.K. BHATTACHARJEE (1980): Orchids of India, Naya Prakash, Calcutta, India. pp. 148, 472.
- BRUHL, P. (1926): A Guide to the Orchids of Sikkim. Calcutta and Simla, Thacker, Spink & Co. pp. 153-138.
- GRANT, C.B. (1895): Orchids of Burma (including the Andaman Islands). Hanthawady Press, Rangoon, p. 311.
- HOOKE, J.D. (1890): Flora of British India. 6: 58, 72, Reeve & Co., Kent, England.
- HUNT, P.F. & M.A. GRIERSON (1973): Orchidaceae. The Bourton Press, London. p. 25.
- JAYAWEERA, D.M.A. (1981): Orchidaceae. In: Dassanayake, M.D. and F.R. Fosberg (eds). A revised handbook to the flora of Ceylon 2. Smithsonian Inst., and the Nat. Sci. Foundation, Washington, D.C. by Amerind Publishing Co. Pvt. Ltd. New Delhi, India. pp. 251-258.
- PRADHAN, U.C. (1979): Indian Orchids: Guide to Identification & Culture. 2: 552-554, 631. U.C. Pradhan, Rishi Road, Kalimpong, India.
- SEIDENFADAN, G. (1972): Contrb. Orch. Fl. Thai. 4. Bot. Tidsskr. 67.
- SMITH, J.J. (1912): Bull. Jard. Bot. Buitenzorg 2. Ser. 8.

34. *BOLBOSCHOENUS PLANICULMIS* (F. SCHMIDT) T. KOYAMA
(CYPERACEAE) — A NEW RECORD FOR SOUTH ASIA
FROM GUJARAT COAST

(With two text-figures)

While working on Cyperaceae specimens from the West Coast of the country, two interesting specimens were found labelled as *Scirpus maritimus* L. var. *affinis* Clarke. As some striking, superficial differences were found in these specimens, a detailed study was made and the specimens were subsequently identified as *Bolboschoenus planiculmis* (Schmidt) Koyama which is not yet reported from India and the neighbouring countries, except from the Chinese coast. A detailed description of the species along with illustrations and distribution is given below.

Bolboschoenus planiculmis (F. Schmidt) T. Koyama, Fl. Taiwan 5: 207. 1978. *Scirpus planiculmis* F. Schmidt, Reisen Amerlande U. Ins. Sachalin 190, t. 8, f. 1-7. 1868; Koyama in Journ. Fac. Sci. Univ. Tokyo 3, 7(6): 330, f. 13. 1958. Fig. 1.

Perennials with long stolons, 30-50 cm high, stramineous when dry. Stolons slender, 1-1.5 mm thick, covered by brownish sheaths, ending in small tubers; tubers ovoid to oblong-ellipsoid, 10-12 x ca 3 mm, rooting and producing new stems. Stems solitary from the




Fig. 1: *Bolboschoenus planiculmis* (Schmidt) Koyama (A) Habit; (B) Spikelet; (C) Glume; (D) Flower in later stage; (E) Stamen; (F) Nut.

tuberous base, narrow, erect, trigonous with flat sides, 1-2 mm thick in the middle portion, nodose towards base, striate, glabrous, few-leaved in the basal half. Leaves cauline, shorter than stem, triangular, almost 3-winged towards apex, linear, acute at apex, 1.5-2 mm wide; basal leaves almost bladeless; sheaths tightly closed, truncate at mouth; ligule absent. Inflorescence a single pseudolateral spikelet. Involucral bracts 1 or 2; main bract erect, as if continuation of the stem, trigonous, 7-15 cm long; smaller one 1.5-4.5 cm long, flat. Spikelets ovoid to broadly ovoid, subacute at apex, 8-20 x 6-10 mm, ferrugineous, densely flowered. Glumes scarious, broadly ovate, emarginate at apex, *ca* 8 x 4-6 mm, puberulous in the upper half; midrib very prominent, extending to a long awn; side veins 0-4. Perianth bristles 4-6, slender, unequal, shorter than or overtopping the nut, retrorsely scabrous.

Stamens 3; filaments flat, hyaline, elongate up to 8 mm; anthers linear, oblong-lanceolate, with a prominent bristly connective appendage at apex, 4-5 mm long. Style *ca* 4 mm long, slightly narrowed to base; stigmas 2, 2.5-3 mm

long. Nuts planoconvex, broadly obovate, cuneate to base, apiculate at apex, 3.5-4.5 x 2.5-3 mm, smooth, creamish-brown to blackish brown; epidermal cells isodiametric, usually not clearly visible in mature nuts.

Fl. & Fr.: October

Habitat: Seashore.

Distribution: So far reported in East and Far East Asia, from the coasts of China, Taiwan, Korea, Japan (including Ryukyus Archipelago) and Sakhalin. Therefore, its present report from south Asia (Gujarat coast in western India) is phytogeographically interesting.

Specimen examined: Aliahet-Hansot, Gujarat, *Toor* 25817. 18.x.1957 (BSI).

According to Koyama (*loc. cit.*), *Bolboschoenus planiculmis* varies in the number of spikelets and also in the shape and size of the nuts, though it is clearly distinct from the closely related, *B. maritimus* complex especially to the subspecies *affinis* (Roth) Koyama. The differences between these two taxa are based on the Indian specimens available in BSI.

B. planiculmis can be distinguished easily by its pseudolateral inflorescence of a single

B. maritimus ssp. *affinis* (Roth) Koyama

Stolons thick and woody
 Tubers stout, 10-25 x 6-9 mm.
 Stems 2.5-6 mm thick near the base (including the sheaths).
 Leaves flat, 2.5-5 mm wide.
 Inflorescence a terminal cluster of 2-6 sessile spikelets, at times reduced to a single spikelet.
 Involucral bracts 2-3, flat, foliaceous.
 Glumes oblong-lanceolate, 7-10 x 2.5-3 mm; side veins usually absent.
 Perianth bristles shorter than the nut.
 Staminal filaments elongate up to 6 mm; anthers oblong, *ca* 2 mm long.
 Style *ca* 2 mm long.
 Nuts *ca* 2.5 x 2 mm.

B. planiculmis (Schmidt) Koyama

Stolons slender, not woody
 Tubers thin, 10-12 x *ca* 3 mm.
 Stems 1.5-3 mm thick near the base (including the sheaths).
 Leaves triangular, 1.5-2 mm wide.
 Inflorescence a single, pseudolateral spikelet.
 Involucral bract 1 or 2; the main bract trigonous, looks like continuation of the stem.
 Glumes broadly ovate, *ca* 8 x 4-6 mm; side veins 0-4.
 Perianth bristles shorter than or overtopping the nut.
 Staminal filaments elongate up to 8 mm; anthers linear or oblong-lanceolate, 4-5 mm long.
 Style *ca* 4 mm long.
 Nuts 3.5-4.5 x 2.5-3 mm.

MISCELLANEOUS NOTES

spikelet and triangular nature of the leaves and the main bract. Slender stolons which are not woody, also characterise the species. Moreover, it is found only along seashores.

Indian Liaison Officer at Royal Botanic Gardens, Kew for literature.

March 18, 1998

V.P. PRASAD
N.P. SINGH

*Botanical Survey of India,
Western Circle, Pune 411 001,
Maharashtra,
India.*

ACKNOWLEDGEMENTS

We thank Dr. P.K. Hajra, Director, Botanical Survey of India, Calcutta for facilities and encouragement and Dr. Sri Krishna Murti,

